2,4-D Sorption in Iron Oxide-Rich Soils: Role of Soil Phosphate and Exchangeable Al

DHARNI VASUDEVAN* AND
ELLEN M. COOPER

Nicholas School of the Environment and Earth Sciences, Duke University, Box 90328, Durham, North Carolina 27701-0328

This study examined herbicide retention in iron oxide-rich variable charge soils (Ultisols) under no cultivation (forest), agriculture (farm), and turf maintenance (golf course) to explore the following hypothesis: inorganic phosphate accumulation from soil fertilization and liming to decrease exchangeable aluminum (Al) content will influence carboxylic acid herbicide sorption onto soils and leaching into groundwater. A suite of soil properties, including mineralogy (particularly soil iron and aluminum oxide content), exchangeable Al content, and soil phosphate content, influenced sorption of the anionic, 2,4-D. In general, 2,4-D sorption was lower in the presence of phosphate,

Methodology

Solutes and Soils. 2,4-D (2,4-dichlorophenoxyacetic acid; Aldrich), the test carboxylic acid herbicide, has a pK_a of 2.5 - 2.7 (24) and is predominantly anionic above pH 3.5 (Figure 1). Alachlor (2-chloro-2′,6′-diethyl-N-(methoxymethyl)acetanilide; Chemservice), the test nonionic (neutral) herbicide, does not have a carboxylate functionality or other groups capable of interacting with surficial Fe and Al atoms. Additionally, the absence of charge in alachlor precludes its electrostatic interaction with charged surfaces.

Three Ultisols of the Georgeville series (fine, kaolinitic, and Typic Kanhapludults), each representing distinct land uses (farm, forest, and golf course) and hence distinct P fertilization and liming histories, were obtained from the North Carolina Piedmont in the United States. All soils had no prior history of 2,4-D or alachlor application. A total of nine unaltered soil samples (3 depth samples from each site)