

INTRODUCTION

Previous studies have suggested that populations of North American aerial avian insectivores are decreasing (Robbins et al. 1989, Böhning-Gaese et al. 1993, Askins 1995). Most recently, based on North American Breeding Bird Survey (BBS) data from between 1966 and 2006 and including most BBS routes for Canada and the United States (excluding Alaska, Yukon, and states or provinces with 15 or fewer BBS routes), Nebel et al. (2011) concluded that Tree Swallow (*Tachycineta bicolor*) populations were among the species of decreasing aerial insectivores, especially in northeastern North America. BBS data, such as those analyzed by Nebel et al. (2011), are collected along roads, and are most concentrated near areas of high human population density (see Sauer et al. 2011 for details on spatiotemporal coverage). In many parts of theaTt

Fig. 1. Location and occupancy rates over time of Tree Swallow sites. Scatterplots associated with open circles over sites have negative trends (Table 1); closed circles indicate increasing trends.

Table 1. Site details (Fig. 1) and results of logistic regressions with number of occupied boxes and number of boxes as response variables, year as explanatory variable, and error distributions set as quasi-binomial. Sites are arranged by longitude. β is the slope, and SE β is the standard error of the slope estimate. Significant slopes in bold.

Site name	Site name, researchers	Latitude
-----------	------------------------	----------

RESULTS

Ten of 16 sites had significant increases or decreases in box occupancy rates (Fig. 1, Table 1). All six sites east of -78° W longitude had decreased occupancy (5 of these significantly so), whereas only two of ten sites west of -78° W longitude had decreased occupancy. Both North ($F_{1,13} = 10.9$, $P = 0.006$) and West ($F_{1,13} = 11.2$, $P = 0.005$) were significant predictors of trend slopes (overall model $R^2 = 0.47$, $F_{2,13} = 5.7$, $P = 0.02$). If the Saskatchewan, Manitoba, and Wisconsin sites were excluded because they regularly had $\sim 100\%$ occupancy, North ($F_{1,10} = 8.4$, $P = 0.02$) and West ($F_{1,10} = 8.6$, $P = 0.01$) still remained significant predictors of trend slopes (overall model $R^2 = 0.47$, $F_{2,10} = 4.4$, $P = 0.04$).

We reanalyzed data from Manitoba after excluding 2005, a year that followed two consecutive years of 100% predation by short-tailed weasels (*Mustela erminea*) (Fig. 1); this did not produce a significant trend for this site ($\beta = 0.041 \pm 0.125$, $t = 0.3$, $P = 0.75$). Excluding 2008 from the Michigan data following a year of complete predation by cats (*Felis domesticus*) and raccoons (*Procyon lotor*; Fig. 1) resulted in a statistically significant positive trend for this site ($\beta = 0.095$

to be behaviorally cryptic (Smith 1978, Stutchbury and Robertson 1985, Shutler and Weatherhead 1991, 1992, 1994).

routes, or timing. Our results indicate that populations of Tree Swallows bear close monitoring; they may provide keys to identifying many of the unknowns contributing to changes in populations of other species of aerial insectivores.

Responses to this article can be read online at:
<http://www.ace-eco.org/vol7/iss1/art3/responses/>

Acknowledgments:

With exception of the first author, author order was determined by the number of box-years contributed to the dataset. We thank Phil Taylor for suggesting this endeavor and various forms of assistance, Megan Colwell, Holly Lightfoot, and Adele Mullie for feedback, legions of students and colleagues who have helped us collect these data, and two reviewers for extraordinarily thorough comments. We acknowledge the enormous effort of volunteers and government agencies in providing BBS data and access to it. Funding sources included NSF (in Ithaca by NSF grants IBN-013437, IBN-920723, IBN-0131437, DEB-0717021 to D. W. Winkler, and IOS-0744753 to Winkler and C. Vleck) and NSERC. The Ontario-Long Point site was funded continuously since its inception by Long Point Bird Observatory (a program of Bird Studies Canada after 1997). For the sake of brevity, we refer readers to our other Tree Swallow publications for acknowledgments of research assistants and additional funding sources.

- Systematics 13:1-21. <http://dx.doi.org/10.1146/annurev.es.13.110182.000245>
- Hussell, D. J. T. 1983. Tree Swallows raise two broods in a season. *Wilson Bulletin* 95: 470-471.
- Hussell, D. J. T. 2003a. Two more double-brooded Tree Swallows. *North American Bird Bander* 28: 49-51.
- Hussell, D. J. T. 2003b. Climate change, spring temperatures, and timing of breeding of Tree Swallows (*Tachycineta bicolor*) in southern Ontario. *Auk* 120:607-618. [http://dx.doi.org/10.1642/0004-8038\(2003\)120\[0607:CCSTAT\]2.0.CO;2](http://dx.doi.org/10.1642/0004-8038(2003)120[0607:CCSTAT]2.0.CO;2)
- Hussell, D. J. T. 2012. The influence of food abundance on nest box occupancy and territory size in the Tree Swallow, a species that does not defend a feeding territory. *Condor* in press.
- Hussell, D. J. T., M. H. Mather, and P. H. Sinclair. 1992. Trends in numbers of tropical and temperate-wintering migrant landbirds at Long Point, Ontario, 1961-1988. Pages 101-114 in *Ecology and conservation of neotropical migrant landbirds* (J. M. Hagan III and D. W. Johnston, Eds.). Smithsonian Inst., Washington, D. C., U.S.A.
- Hussell, D. J. T., and T. E. Quinney. 1987. Food abundance and clutch size of Tree Swallows *Tachycineta bicolor*. *Ibis* 129:243-258. <http://dx.doi.org/10.1111/j.1474-919X.1987.tb03204.x>
- Jones, J. 2003. Tree Swallows (*Tachycineta bicolor*): A new model organism? *Auk* 120:591-599. [http://dx.doi.org/10.1642/0004-8038\(2003\)120\[0591:TSTBAN\]2.0.CO;2](http://dx.doi.org/10.1642/0004-8038(2003)120[0591:TSTBAN]2.0.CO;2)
- Keller, C. M. E., and J. T. Scallan. 1999. Potential roadside biases due to habitat changes along Breeding Bird Survey routes. *Condor* 101:50-57. <http://dx.doi.org/10.2307/1370445>
- Kempnaers, B., S. Everding, C. Bishop, P. Boag, and R. J. Robertson. 2001. Extra-pair paternity and the reproductive role of male floaters in the Tree Swallow (*Tachycineta bicolor*). *Behavioral Ecology and Sociobiology* 49:251-259 <http://dx.doi.org/10.1007/s002650000305>
- Lifjeld, J. T., P. O. Dunn, R. J. Robertson, and P. T. Boag. 1993. Extra-pair paternity in monogamous Tree Swallows. *Animal Behaviour* 45:213-229. <http://dx.doi.org/10.1006/anbe.1993.1028>
- Lombardo, M. P. 1988. Evidence of intraspecific brood parasitism in the Tree Swallow. *Wilson Bulletin* 100:126-128.
- Lorch, J. M., C. U. Meteyer, M. J. Behr, J. G. Boyles, P. M. Cryan, A. C. Hicks, A. E. Ballmann, J. T. H. Coleman, D. N. Redell, D. M. Reeder, and D. S. Blehert. 2011. Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature* 480:376-378. <http://dx.doi.org/10.1038/nature10590>
- Monroe, A. P., K. K. Hallinger, R. L. Brasso, and D. A. Cristol. 2008. Occurrence and implications of double brooding in a southern population of Tree Swallows. *Condor* 110:382-386. <http://dx.doi.org/10.1525/cond.2008.8341>
- Muldal, A., H. L. Gibbs, and R. J. Robertson. 1985. Preferred nest spacing of an obligate cavity-nesting bird, the Tree Swallow. *Condor* 87:356-363. <http://dx.doi.org/10.2307/1367216>
- Nebel, S., A. Mills, J. D. McCracken, and P. D. Taylor. 2011. Declines of aerial insectivores in North America follow a geographic gradient. *Conservation and Ecology - Écologie et conservation des oiseaux* 5(2):1. [online] URL: <http://www.ace-eco.org/vol5/iss2/art1/>.
- Pabian, S. E., and M. C. Brittingham. 2011. Soil calcium availability limits forest songbird productivity and density. *Auk* 128:441-447. <http://dx.doi.org/10.1525/auk.2011.10283>
- Paynter, R. A. 1954. Interrelations between clutch-size, brood-size, pre fledging survival, and weight in Kent Island Tree Swallows. *Bird Banding* 25: 35-38, 102-110, 136-148.
- Pulliam, H. R. 1988. Sources, sinks, and population regulation. *American Naturalist* 132:652-661. <http://dx.doi.org/10.1086/284880>
- R Core Development Team. 2006. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Robbins, C. S., J. R. Sauer, R. S. Greenberg, and S. Droege. 1989. Population declines in North American birds that migrate to the neotropics. *Proceedings of the National Academy of Science (U.S.A.)* 86:7658-7662. <http://dx.doi.org/10.1073/pnas.86.19.7658>
- Robertson, R. J., and W. B. Rendell. 1990. A comparison of the breeding ecology of a secondary cavity nesting bird, the Tree Swallow (*Tachycineta bicolor*), breeding in nest-boxes and natural cavities. *Canadian Journal of Zoology* 68:1046-1052. <http://dx.doi.org/10.1139/z90-152>
- Robertson, R. J., and B. J. Stutchbury. 1988. Experimental evidence for sexually selected infanticide in Tree Swallows. *Animal Behaviour* 36:749-753 [http://dx.doi.org/10.1016/S0003-3472\(88\)80158-1](http://dx.doi.org/10.1016/S0003-3472(88)80158-1)
- Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2011. The North American Breeding Bird Survey, Results and Analysis 1966 - 2009. Version 3.23.2011. U.S.G.S. Patuxent Wildlife Research

