

Host Plant Preferences of *Philaenus spumarius*, the meadow spittlebug

Zoe Wood, Class of 2018

I had the exciting opportunity this summer to work with a system that no one had previously examined in depth on Kent Island: *Philaenus spumarius*, the meadow spittlebug. While Kent is a wonderful place to study seabirds and migrating birds, it also contains a wealth of insect and plant life that make great subjects.

Spittlebugs are small but visible crop pests that are easily recognized by the frothy spittle masses the nymph creates on the plant it feeds on. Spittlebugs are generalist xylem-sap feeders, meaning they are not host-plant specific to any one species but rather are versatile in their feeding behavior. Nevertheless, they still have preferences for certain plants, including but not limited to nitrogen-fixing angiosperms, suggesting the role of nitrogen content as a potential factor in host plant selection¹. Plants with higher nitrogen content may consequently be a richer source of amino acids for the xylem-feeding insects. My project aimed to characterize the 1) diversity of the plants on Kent found to be spittlebug hosts, 2) density and performance of nymphs on different plant species, 3) and changes in density on goldenrod (*Solidago rugosa*) over the course of eight weeks with nitrogen fertilizer application.

So far, I've been able to identify 32 host plants and trees species, including several flowering plants and some conifers. I measured nymph body sizes on some of these plants as a proxy for performance, and found that on average, nymphs on goldenrod were larger than nymphs on sheep sorrel. For my density censuses, I created paired goldenrod plots (half fertiliz