

Explore the space: the behavioral effects of semaphorin-induced neuroplasticity in the nervous system of *Gryllus bimaculatus*
Ean L. Small, Class of 2023

Scholars have long examined the phonotactic behavioral response of crickets induced by sudden neurological changes resulting from sound stimuli. This research has led neuroethologists (scientists studying animal behavior) to an intensive exploration of the flying functions of the cricket auditory system. The relevant literature has elucidated the two integral functions of crickets' acoustic response necessary for their survival: sexual partner identification and avoidance of predators (Huber et al., 1989). The predator-prey interactions between bats and crickets have shaped the evolutionary adaptations forming the unique auditory system of crickets. Observational studies have concluded that crickets typically initiate flying activity at night and are preyed upon by bats (Ulagaraj, 1975; Jacobs & Bastian, 2016). While hunting, bats utilize echolocation via high-frequency ultrasonic sounds to identify the location of crickets in complete darkness (Jacobs & Bastian, 2016). In response to the high-frequency pulses generated by bats, crickets turn away from the sound stimuli and attempt to fly away from predators (Moiseff et al., 1978). This behavioral response is known to be negative phonotaxis.

To produce effective behavior, the cricket auditory system displays neuronal regeneration in adults that allows them to recover their behavioral response post-injury. Prior to injury, cricket auditory neurons are confined to one side of the prothoracic ganglia (PTG) midline, sending auditory signals from one side of the body to the same side of the brain. Crickets that are deafened by removing the foreleg containing the cricket auditory organ lose synaptic connections that allow for the relay of signals from the ear to the brain (Horch et al., 2011). This neural degeneration significantly impairs the Horch lab has identified the semaphorin 1a.2 (sema1a.2) protein as a possible molecule involved in the regrowth of neurons across the midline post-injury, which in turn helps crickets recover their behavioral response.

This project sought to correlate the axon guidance nature of sema1a.2 with a unique cricket behavioral response to predatorial sound stimuli. Over the course of this summer, I worked to analyze extensive past data obtained in the Horch Lab. Crickets analyzed in this data were larval stage crickets injected with double-stranded RNA.

Literature Cited

- Jacobs DS, Bastian A. Chapter 2; Bat Echolocation: Adaptations for Prey Detection and Capture. In: Predator-prey interactions: Coevolution between bats and their prey. Cham: Springer International Publishing; 2016. p. 129.
- Horch, H.W., et al. Bilateral consequences of chronic unilateral deafferentation in the auditory system of the cricket *Gryllus bimaculatus*. *Dev Neurosci*, 2011;33(1): p. 2137.
- Huber F, M.T., Loher W (eds). Cricket behavior and neurobiology. Comstock Pub Associates, Ithaca, 1989.
- Moiseff A, Pollack GS, Hoy RR. Steering responses of flying crickets to sound and ultrasound: Mate attraction and predator avoidance. *Proceedings of the National Academy of Sciences*. 1978;75(8):4052-4056. doi:10.1073/pnas.75.8.4052
- Ulagaraj SM. Mole crickets: Ecology, behavior, and dispersal flight (Orthoptera: Gryllotalpidae: Scapteriscus). *Environmental Entomology*. 1975;4(2):267-273. doi:10.1093/ee/4.2.265